




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Evaluation of Invasive Ventilation Rate and Comorbidities, Clinical Signs and Lab Findings among COVID 19 Patients

Nina Farzan, Sepideh Vahabi, Behrooz Farzan, Shima Sadat Hashemi Madani



PII: S2052-2975(21)00089-5

DOI: <https://doi.org/10.1016/j.nmni.2021.100925>

Reference: NMNI 100925

To appear in: *New Microbes and New Infections*

Received Date: 13 November 2020

Revised Date: 21 April 2021

Accepted Date: 19 July 2021

Please cite this article as: Farzan N, Vahabi S, Farzan B, Hashemi Madani SS, Evaluation of Invasive Ventilation Rate and Comorbidities, Clinical Signs and Lab Findings among COVID 19 Patients, *New Microbes and New Infections*, <https://doi.org/10.1016/j.nmni.2021.100925>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 The Author(s). Published by Elsevier Ltd.

Research article

Evaluation of Invasive Ventilation Rate and Comorbidities, Clinical Signs and Lab Findings among COVID 19 Patients

Running title: Invasive Ventilation Rate among COVID 19 Patients

Nina Farzan¹, Sepideh Vahabi², Behrooz Farzan^{2*}, Shima Sadat Hashemi Madani³

1-Department of Emergency Medicine, Clinical Research development center, Qom university of medical sciences, Qom, Iran

2-Department of Anesthesiology, Faculty of Medicine, Lorestan university of Medical Sciences, Khorramabad, Iran.

3-Emergency Medicine Department, Student Research development center, Qom university of medical sciences, Qom, Iran

Corresponding author: Behrooz Farzan, Assistant professor of Anesthesiology, Lorestan university of Medical Sciences, Khorramabad, Iran.

Conflict of interest: The authors deny any conflict of interest in any terms or by any means during the study.

Ethical approval and consent to participate

All procedures performed in this study involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Consent for publication: Not applicable

Availability of data and material: Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

Funding Source: No funding was secured for this study.

Authors Contributions

Dr. Nina Farzan: Planned the study, wrote the protocol, collected the data and drafted the manuscript and accepted the final draft.

Dr. Sepideh Vahabi: Planned and designed the study, collected the data.

Dr. Behrooz Farzan: analyzed the data and critically revised the draft and finally approved the manuscript.

Abstract

The severity of COVID-19 has been to be associated with the comorbidities. It is defined as the presentation of severe respiratory dysfunction or failure, leading to the need of ventilation and mortality. The aim of this study is the evaluate the factors predicting the rate of invasive ventilation among these patients.

This retrospective study involved 317 COVID-19 patients referred to (XXX) Hospital in Qom, Iran. The following data was obtained for all the patients: demographic parameters, comorbidities, need of mechanical ventilation, signs and symptoms and lab findings.

The results from the demographic data of the study indicated that the need of mechanical ventilation is significantly associated with advanced age, $p=0.001$. Additionally, hypertension, leukopenia and blood urea nitrogen to creatinine ratio $p=0.008$. $p=0.042$ and $p<0.001$, respectively are significantly associated with increased need of mechanical ventilation. Malignancy, diabetes, asthma, chronic obstructive pulmonary disease, headache, fever, platelet count, prothrombin time, c-reactive protein, erythrocyte sedimentation rate and creatinine phosphatase were not significantly different in the two groups, $p>0.05$.

Prediction of the extent of severity among COVID-19 patients using clinical parameters and comorbidities prepare medical practitioners and health care centers to take immediate measurements and reduce the burden of scarcity of health supplies and care.

Keywords: COVID-19, comorbidities, invasive ventilation.

Introduction

The novel pneumonia-causing coronavirus (2019-nCoV) leading to coronavirus disease 2019 (COVID-19) is a single-stranded RNA virus and is known to be close to SARS virus (severe acute respiratory syndrome virus). The first case of COVID-19 was reported in Hubei, China where later it was considered as an epidemic. In March 2020, the cases of COVID-19 were 13 times more in the other regions of the world and it was declared as the pandemic by the World Health Organization (WHO) (1). According to the WHO report published on 13th December, 2020, the overall number of global cases are 70,476,836 and 1,599,922 deaths have been reported (2).

The real-time polymerase chain reaction (RT-PCR) using nasal throat sample is used for the diagnosis of the infection. Additionally, chest CT (computed tomography) scan is performed to determine the extent of lung involvement (3). The common signs and symptoms of the disease include fever, dry cough, headache, fatigue dyspnea and vomiting (4). Severe cases of the infection can lead to organ failure and death while ordinary respiratory distress is common in non-severe patients (5). The presentation of respiratory distress is seen after 7 days of infection. Patients are likely to develop silent hypoxemia as well (hypoxemia without respiratory distress)(6). Several biomarkers can predict the severity of the disease and might be helpful in determining the need of critical care in patients(7). Increased c-reactive protein and lymphopenia are likely to be good predictors of hypoxemia and associated mortality (8, 9). The incidence of COVID-19 among patients with comorbidities is 30-50%. Hypertension and respiratory and cardiovascular disorders can add to the severity of the disease (10). The treatment of the disease using remdesivir and other antiviral is under clinical trials (NCT04252664 and NCT04257656) (8).

The aim of this study is to evaluate the incidence of comorbidities and sign and symptoms of COVID-19 in the prognosis of invasive ventilation.

Methods:

This retrospective study included COVID-19 adult patients who underwent invasive ventilation at (XXX). Patients aged ≥ 18 years diagnosed with COVID-19 according to the WHO guideline from February 2020 - March 2020 referred to our center were included.

The information regarding demographic characteristics, clinical symptoms, lab findings, treatment and the need of invasive mechanical ventilation were collected from the record of these patients and were entered in a standardized data form. The treatment of the patients was provided according to the guideline of Infectious Diseases Society of America Guidelines on the Treatment and Management of COVID-19 Infection (11) and Iranian Expert's Consensus Statement (12). Additionally, mild, moderate and severe cases were also determined based on these guidelines, that also determined the need of invasive mechanical ventilation in these patients. 7.0- to 8.0-mm endotracheal tube (EET) was placed between vocal cords into the trachea as per standard procedure of invasive ventilation.

The diagnosis of COVID-19 in symptomatic patients was made using RT-PCR using throat-swab specimen. The routine lab tests included complete blood count, renal and liver function tests, coagulation profile and c-reactive protein (CRP). The CT scan was also performed in all the patients.

The data obtained was digitized and statistically analyzed using SPSSv22. The mean and variance indices were used to describe the concentration and distribution of quantitative data and relative frequency and frequency indices for qualitative variables. Fisher's chi-square and precision tests were used to compare the rate of invasive ventilation in qualitative variables. P value < 0.05 was considered to be statistically significant.

Results:

A total of 317 patients were included in the study where 167 were male and 139 were female. The average age of patients was 59.71 ± 16.46 years. Advanced age was significantly associated with the need of invasive ventilation, $p=0.001$ (Figure1). The need of invasive ventilation was not different among the two genders, $p=0.44$.

The rate of arthralgia was 30.2% for patients with invasive ventilation and 39.1% for patients without invasive ventilation. The chi-square test showed no significant relationship between arthralgia and invasive ventilation, $p=0.193$.

Among asthmatic COVID-19 patients, the rate of invasive ventilation was 3.2% and non-asthmatic patients was 4.6%. Fisher's exact test showed no significant relationship between asthma and invasive ventilation, $p=0.465$.

The rate invasive ventilation among patients with malignancy was 4.8% and among the patients without malignancy was 2.1%. The results from Fisher's exact test showed that there was no significant relationship between malignancy and invasive ventilation, $p = 0.371$.

The rate of invasive ventilation among patients with BUN to creatinine ratio greater than 20:1 was 42.9% and 11.8% for patients without invasive ventilation. The Chi-square test showed a significant association between BUN to creatinine ratio and invasive ventilation, $p<0.001$ (Figure2).

The COPD rate was 4.8% among patients needing invasive ventilation and 1.7% in patients without invasive ventilation. The Fisher's exact test showed that there was no significant relationship between the incidence of COPD and the need of invasive ventilation, $p=0.162$.

The CPK level was high in 3.2% patients with invasive ventilation and 2.5% patients without invasive ventilation. The Fischer's exact test showed that the levels of CPK was not significantly associated with the need of invasive ventilation, $p = 0.675$.

The need of invasive ventilation among patient with CRP > 3.0 mg/L was 46% and 52.9% for patients without invasive ventilation. The findings from the Chi-square test showed that the levels of CRP were not significantly different among patients with or without invasive ventilation $p=0.675$.

The need of invasive ventilation among DM patients was 23.8% and 26.1% DM patients did not need invasive ventilation. The Chi-square test showed no significant relationship between the prevalence of DM and the need of invasive ventilation, $p=0.717$.

The incidence of olfactory dysfunction was 4.8% in patients with invasive ventilation and 3.4% in patients without invasive ventilation. The Fisher's exact test showed that no significant relationship between olfactory dysfunction and invasive ventilation, $p=0.704$.

The incidence of anorexia nervosa was 6.3% among patients with invasive ventilation and 5.9% in patients without invasive ventilation. The Fisher's accurate test showed no significant relationship between anorexia nervosa and invasive ventilation, $p=1$.

The invasive ventilation was provided to 1.6% patients with diarrhea and 0.8% non-diarrhea patients. The results from the Fisher's exact tests did not show any significant relationship between the incidence of diarrhea and the need of mechanical ventilation, $p=0.507$.

The incidence of increased ESR (1–13 mm/hr for males and 1–20 mm/hr for females) was 28.6% for patients with invasive ventilation and 26.1% for patients without invasive ventilation. The Fisher's exact test showed that ESR was not correlated with the need of invasive ventilation among COVID-19 patients, $p=0.678$.

The incidence of hypertension was 37.7% among patients with invasive ventilation and 23.1% in patients without invasive ventilation. The Chi-square test showed a significant association

between the incidence of hypertension and invasive ventilation, $p=0.008$. The need of mechanical ventilation was 55% higher among hypertensive patients(Figure3).

The incidence of fatigue was 12.7% among patients receiving invasive ventilation and 12.6% in patients without invasive ventilation. The Chi-square test showed no significant relationship between fatigue and invasive ventilation, $p=1$.

The incidence of IDH was 23.2% among invasive ventilation patients and 16.4% in patients without invasive ventilation. The Chi-square test showed no significant relationship between IDH and invasive ventilation, $p=0.172$.

The incidence of lymphopenia was 42.9% in patients with invasive ventilation and 29.4% in patients without invasive ventilation. The results from Chi-square test showed that the incidence of lymphopenia was significantly greater among patients needing invasive ventilation, $p=0.042$ (Figure4). The risk of COVID-19 patients with lymphopenia needing invasive ventilation was 1.79 times higher than patients without lymphopenia.

The incidence of leukopenia was 7.9% in patients receiving invasive ventilation and 17.4% in patients without invasive ventilation. The Chi-square test showed no significant relationship between leukopenia and invasive ventilation, $p=0.068$.

The rate of invasive ventilation among transplant recipients and non-recipient was 3.2% and 0.8%, respectively. The Chi-square showed no significant difference in the need to ventilation among these two groups $p=0.194$.

1.6% patients with platelet count lesser than 150,000 received invasive ventilation and 7.6% for patients did not receive invasive ventilation. The Fisher's exact test showed no significant relationship between platelet count and the need of ventilation $p=0.14$.

For 1.6% with prothrombin time ≤ 16 seconds, invasive ventilation was provided whereas, 1.3% of these patients did not receive invasive ventilation. The Fisher's exact test showed no significant difference between the two groups, $p= 1$.

The incidence of headache among patients needing and not needing invasive ventilation was 4.8% and 9.2%, respectively. The Chi-square did not show any significant difference among the two groups, $p=0.252$.

The rate of cough among patients needing and those not needing mechanical ventilation was 60.3% and 58.4%, respectively. The Chi-square test did not show any significance difference among the two groups, $p=0.784$.

The rate of productive cough was 9.5% in patients with invasive ventilation and 7.1% among patients without invasive ventilation. The Fisher's exact test showed that there was no significant relationship between productive cough and invasive ventilation, $p= 0.953$.

The need of mechanical ventilation among patients with fever was 71.4%. 69.3% of pyretic patients did not need mechanical ventilation. The Chi-square test showed no significant association between fever and invasive ventilation, $p=0.747$.

The rate of nausea was 12.7% in patients with invasive ventilation and 10.5% among patients without invasive ventilation. The Chi-square test showed no significant relationship between nausea and invasive ventilation, $p=0.620$.

Respiratory distress was 87.3% among patients with invasive ventilation and 87.8% in patients without invasive ventilation. The Chi-square test showed no significant relationship between shortness of breath and invasive ventilation, $p = 0.860$.

Discussion

Acute hypoxemic respiratory insufficiency/failure is one of the common presentations of COVID-19, needing ventilation and intubation(13, 14). The severity of the disease is defined as tachypnoea ≥ 30 breaths per min, oxygen saturation $\leq 93\%$ at rest, or $\text{PaO}_2/\text{FiO}_2$ ratio < 300 mm Hg and respiratory failure requiring mechanical ventilation, septic shock, or other organ dysfunction or failure, requiring intensive care (15). Our study evaluated the relationship between the need to mechanical ventilation with commodities and clinical and lab presentation.

Goyal P et al (16) reported that obesity, male gender, elevated liver function test, inflammatory markers are predictors of invasive ventilation in COVID-19 patients. Our study did not report any significantly difference among the levels of inflammatory markers like CRP and ESR in invasive ventilation and non-invasive ventilation group. The patients presented with hypertension, cardiovascular disease and diabetes mellitus are at the greater risk of the infection. 2019-nCoV binds to the angiotensin converting enzyme-2, which is widely expressed in lung epithelium. Its

expression is elevated in diabetic and hypertensive patient, which can increase the risk of infection in these patients. The result of our retrospective study revealed that the need of invasive mechanical ventilation is significantly greater in hypertensive patients however, it was untrue for diabetic patients. These outcomes indicate that hypertension is likely to be associated with the severity of the disease marked with severe respiratory distress (17).

Eosinophilia and early onset of shortness of breath are also associated with increased risk of mortality (18). A single center retrospective study by Chen Q et al (19) on 145 patients reported that the severity of the disease is significantly associated with advanced age, diabetes mellitus, anorexia, chest tightness, greater body mass index, and dyspnea. The outcomes of our study indicated that these parameters do not identify the potential candidates for invasive ventilation.

COVID-19 severity (defined by ICU admission, need of mechanical ventilation or mortality) is significantly associated with low-platelet count (thrombocytopenia). Morphological change in the capillary bed of the lung epithelial can affect platelet defragmentation. It is also hypothesized that coronavirus can affect bone marrow leading to abnormal hematopoiesis (20). Furthermore, coagulopathy and prothrombin time ≥ 16 is also significantly associated with in-hospital death (21). Nonetheless, platelet count and prothrombin time were not significantly associated with the need of invasive ventilation in our patient group.

Zhang L et al (22) reported that the severity of the disease among cancer patients infected with coronavirus 2019 is associated with a greater need of invasive mechanical ventilation. The need of mechanical ventilation among these patients were recorded to be 35.7%. In our study group, the requirement of invasive mechanical ventilation was not associated with the presence of malignancy. The outcomes are likely to dependent on the stage of cancer and the usage of immunosuppressant drugs.

The outcomes from a very limited number of renal and heart transplant patients have suggested that transplant recipients are likely to present the similar mild COVID-19 symptoms as non-transplant patients due to the modulation of inflammatory response (23, 24). The findings of our study also indicated that the need of invasive ventilation does not differ significantly among transplant and nontransplant recipients.

A meta-analysis and systematic review by Zhao et al (25) concluded that the incidence of COPD is significantly associated with an increased risk of severe COVID-19 and the need of mechanical ventilation. The results of our study are not in parallel with these findings.

Elevated creatine and BUN in kidney disease COVID-19 patients are independent risk factors of in-hospital mortality (26). A study from Cao (27) stated that BUN and creatinine might not be involved in the prognosis of the disease. The results of our study report a significant difference in BUN to creatinine ration among patient who received invasive ventilation and those who did not.

Conclusion:

The continuous increase in the number of COVID-19 cases have led to extensive demand of medical supplies and facilities. Critically ill patients require ICU care and ventilation. Predictors of the need of mechanical ventilation can prepare health care workers and hospital to provide a strategic plan to meet the demands. Further studies involving outcomes from multi centers and a greater number of patients are therefore required.

References

1. Cascella M, Rajnik M, Cuomo A, Dulebohn SC, Di Napoli R. Features, evaluation and treatment coronavirus (COVID-19). Statpearls [internet]: StatPearls Publishing; 2020.
2. Organization WH. Coronavirus disease (COVID-19): weekly epidemiological, update 15. 2020.
3. Wang Y, Wang Y, Chen Y, Qin Q. Unique epidemiological and clinical features of the emerging 2019 novel coronavirus pneumonia (COVID- 19) implicate special control measures. Journal of medical virology. 2020;92(6):568-76.
4. Tian S, Hu N, Lou J, Chen K, Kang X, Xiang Z, et al. Characteristics of COVID-19 infection in Beijing. Journal of Infection. 2020;80(4):401-6.
5. Yang W, Cao Q, Qin L, Wang X, Cheng Z, Pan A, et al. Clinical characteristics and imaging manifestations of the 2019 novel coronavirus disease (COVID-19):A multi-center study in Wenzhou city, Zhejiang, China. Journal of Infection. 2020;80(4):388-93.
6. Alizadeh R, Aghsaiefard Z. Does COVID19 activates previous chronic pain? A case series. Annals of Medicine and Surgery. 2021;61:169-71.
7. Aghsaiefard Z, Alizadeh R. The Role of Angiotensin Converting Enzyme in Immunity: Shedding Light on Experimental Findings. Endocrine, Metabolic & Immune Disorders Drug Targets. 2021.
8. Xie J, Tong Z, Guan X, Du B, Qiu H, Slutsky AS. Critical care crisis and some recommendations during the COVID-19 epidemic in China. Intensive care medicine. 2020:1-4.

9. Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *The Lancet respiratory medicine*. 2020;8(4):420-2.
10. Yang J, Zheng Y, Gou X, Pu K, Chen Z, Guo Q, et al. Prevalence of comorbidities and its effects in patients infected with SARS-CoV-2: a systematic review and meta-analysis. *International Journal of Infectious Diseases*. 2020;94:91-5.
11. Bhimraj A, Morgan RL, Shumaker AH, Lavergne V, Baden L, Cheng V, et al. Infectious Diseases Society of America guidelines on the treatment and management of patients with COVID-19 infection. Website <https://www.idsociety.org/globalassets/idsa/practice-guidelines/covid-19/treatment/idsa-covid-19-gl-tx-and-mgmt-4-11-20-1058-am-edt.pdf> [accessed 12 April 2020]. 2020.
12. Karimi A, Rafiei Tabatabaei S, Rajabnejad M, Pourmoghaddas Z, Rahimi H, Armin S, et al. An Algorithmic Approach to Diagnosis and Treatment of Coronavirus Disease 2019 (COVID-19) in Children: Iranian Expert's Consensus Statement. *Arch Pediatr Infect Dis*. 2020;8(2):e102400.
13. Farzan N, Vahabi S, Madani SSH, Farzan B. Invasive Mechanical Ventilation and Clinical Parameters in COVID19 Patient: Can Age be a Factor? *International Journal of Surgery Open*. 2021:100344.
14. Farzan N, Vahabi S, Shirvani R, Shakeri M, Araghi M, Birjandi M, et al. Evaluation of invasive ventilation (intubation) prognosis in patients with Covid-19 symptoms. *International Journal of Surgery Open*. 2020;27:149-53.
15. Novel CPERE. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. *Zhonghua liu xing bing xue za zhi= Zhonghua liuxingbingxue zazhi*. 2020;41(2):145.
16. Goyal P, Choi JJ, Pinheiro LC, Schenck EJ, Chen R, Jabri A, et al. Clinical Characteristics of Covid-19 in New York City. *New England Journal of Medicine*. 2020.
17. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med*. 2020;8(4):e21-e.
18. Du Y, Tu L, Zhu P, Mu M, Wang R, Yang P, et al. Clinical Features of 85 Fatal Cases of COVID-19 from Wuhan: A Retrospective Observational Study. *American Journal of Respiratory and Critical Care Medicine*. 2020;0(ja):null.
19. Chen Q, Zheng Z, Zhang C, Zhang X, Wu H, Wang J, et al. Clinical characteristics of 145 patients with corona virus disease 2019 (COVID-19) in Taizhou, Zhejiang, China. *Infection*. 2020;1-9.
20. Lippi G, Plebani M, Henry BM. Thrombocytopenia is associated with severe coronavirus disease 2019 (COVID-19) infections: A meta-analysis. *Clinica Chimica Acta*. 2020;506:145-8.
21. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet*. 2020;395(10229):1054-62.
22. Zhang L, Zhu F, Xie L, Wang C, Wang J, Chen R, et al. Clinical characteristics of COVID-19-infected cancer patients: a retrospective case study in three hospitals within Wuhan, China. *Annals of Oncology*. 2020.
23. Siddiqi HK, Mehra MR. COVID-19 illness in native and immunosuppressed states: a clinical-therapeutic staging proposal. *The Journal of Heart and Lung Transplantation*. 2020.


24. Zhu L, Xu X, Ma K, Yang J, Guan H, Chen S, et al. Successful recovery of COVID- 19 pneumonia in a renal transplant recipient with long- term immunosuppression. *American Journal of Transplantation*. 2020.
25. Zhao Q, Meng M, Kumar R, Wu Y, Huang J, Lian N, et al. The impact of COPD and smoking history on the severity of COVID-19: A systemic review and meta-analysis. *Journal of Medical Virology*.n/a(n/a).
26. Cheng Y, Luo R, Wang K, Zhang M, Wang Z, Dong L, et al. Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney International*. 2020;97(5):829-38.
27. Cao W. Clinical features and laboratory inspection of novel coronavirus pneumonia (COVID-19) in Xiangyang, Hubei. *medRxiv*. 2020:2020.02.23.20026963.


Figure1: The relationship between invasive ventilation and age


Figure2: The relationship between invasive ventilation and blood urea nitrogen (BUN) to creatinine ratio.


Figure3: The relationship between invasive ventilation and hypertension

Figure4: The association of invasive ventilation and lymphopenia









Dr. Nina Farzan and Dr.Shima Sadat Hashemi Madani: Planned the study, wrote the protocol, collected the data and drafted the manuscript and accepted the final draft.

Dr. Sepideh Vahabi: Planned and designed the study, collected the data.

Dr. Behrooz Farzan: analyzed the data and critically revised the draft and finally approved the manuscript.